arXiv:1703.09495v1 [math.CA] 28 Mar 2017

A NOTE ON MAXIMAL FOURIER RESTRICTION FOR SPHERES IN ALL DIMENSIONS

MARCO VITTURI

Abstract. We prove a maximal Fourier restriction theorem for the sphere $\mathbb{S}^{d-1}$ in $\mathbb{R}^d$ for any dimension $d \geq 3$ in a restricted range of exponents given by the Stein-Tomas theorem. The proof consists of a simple observation. When $d = 3$ the range corresponds exactly to the full Stein-Tomas one, but is otherwise a proper subset when $d > 3$. We also present an application regarding the Lebesgue points of functions in $\mathcal{F}(L^p)$ when p is sufficiently close to 1.

1. introduction

Very recently a new line of investigation in the field of Fourier restriction studies has been opened by Müller, Ricci and Wright in [2], namely that of maximal Fourier restriction theorems. The goal of such line of investigation is to study the Lebesgue points of the Fourier transform $\widehat{f}$ of a generic function $f \in L^p$ when $1 \leq p \leq 2$ and sufficiently close to 1. In the aforementioned paper they prove that for the case of curves in $\mathbb{R}^2$ the following holds:

Theorem 1.1 ([2])**.** *Let Γ be a C^2 curve in $\mathbb{R}^2$ and let $f \in L^p$ with $1 \leq p < 8/7$. Then, with respect to arclength measure, a.e. point of Γ where the curvature does not vanish is a Lebesgue point of $\widehat{f}$.*

In particular, if $\mathcal{R}$ is the Fourier restriction operator associated to Γ, one has from the above that for all $f \in L^p$ with $1 \leq p < 8/7$

$$\mathcal{R}f = \widehat{f}\big|_\Gamma$$

for a.e. point of Γ where the curvature is non zero (a.e. with respect to arclength measure). Theorem 1.1 is the consequence of a clever trick (which we have included in the proof of Proposition 2.2, for the reader's convenience) and the following

Theorem 1.2 ([2])**.** *Let Γ be the graph of a C^2 function $\gamma : I \to \mathbb{R}$, where I is a bounded interval, and let μ denote the Affine measure on Γ, which for Γ in this form is given by*

$$\mathrm{d}\mu((\xi, \gamma(\xi))) = |\gamma''(\xi)|^{1/3}\mathrm{d}\xi.$$

Let $\chi \in \mathscr{S}(\mathbb{R}^2)$ be a fixed Schwartz function with $\int \chi = 1$ and define the maximal Fourier restriction operator

$$\mathcal{M}f(\xi) := \sup_{\varepsilon,\delta>0} \Big| \int\int_{\mathbb{R}^2} \widehat{f}(\xi + s, \gamma(\xi) + t)\chi_\varepsilon(s)\chi_\delta(t)\mathrm{d}s\mathrm{d}t \Big|.$$

Then the estimate

$$\|\mathcal{M}f\|_{L^q(\Gamma,\mathrm{d}\mu)} \lesssim_{p,q} \|f\|_{L^p(\mathbb{R}^2)} \tag{1.1}$$

holds for all $f \in L^p$ with $1 \leq p < 4/3$ and $p' \geq 3q$.

Date: March 28, 2017.

1991 *Mathematics Subject Classification.* 42B10, 42B25.

Key words and phrases. Fourier restriction, maximal operators.

We have used in the statement the standard notation

$$\chi_\varepsilon(x) = \frac{1}{\varepsilon}\chi(\frac{x}{\varepsilon}).$$

Observe that the range of exponents for which (1.1) holds is the same as that for the usual operator of Fourier restriction to Γ. The proof of the above theorem follows the lines of Sjölin's proof of the Fourier restriction conjecture for curves in the plane as given in [4].

In this short note we consider the case of Fourier restriction to the sphere $\mathbb{S}^{d-1}$ immersed in d-dimensional euclidean space, $d \geq 3$. Let $\mathrm{d}\sigma$ denote the surface measure of the sphere. Define, analogously to the above, the maximal Fourier restriction operator for the sphere

$$\mathscr{M} f(\omega) := \sup_{\varepsilon>0} \Big| \int \widehat{f}(\omega + y)\chi_\varepsilon(y)\mathrm{d}y \Big|,$$

where ω ranges over $\mathbb{S}^{d-1}$ and $\chi \in \mathscr{S}(\mathbb{R}^d)$ is a fixed Schwartz function with $\int \chi = 1$, $\chi_\varepsilon(y) = \varepsilon^{-d}\chi(y/\varepsilon)$. Then we have the following

Theorem 1.3. *Let $d \geq 3$. The operator $\mathscr{M}$ satisfies*

$$\|\mathscr{M} f\|_{L^q(\mathbb{S}^{d-1},\mathrm{d}\sigma)} \lesssim_{p,q,d} \|f\|_{L^p(\mathbb{R}^d)} \tag{1.2}$$

for $1 \leq p \leq 4/3$, $p' \geq \frac{d+1}{d-1}q$.

Moreover, for $1 \leq p \leq 8/7$, if $f \in L^p(\mathbb{R}^d)$ then σ-a.e. point of $\mathbb{S}^{d-1}$ is a Lebesgue point for $\widehat{f}$.

Observe that when $d = 3$ the range of exponents for which (2.1) holds has endpoint $L^{4/3} \to L^2$, and thus corresponds exactly to the Stein-Tomas range for this dimension. For larger values of d, the stated range is however only a subset of the full Stein-Tomas range, which is

$$1 \leq p \leq \frac{2(n+1)}{n+3}, \quad p' \geq \frac{d+1}{d-1}q.$$

It is precisely the fact that the adjoint estimate to $L^{4/3} \to L^q$ is $L^{q'} \to L^4$ that allows for a simple proof of the theorem, since 4 is an even integer and thus the restriction estimate can be restated in cancellation-free form as in (2.3) below. Indeed, one can prove the Stein-Tomas theorem in $d = 3$ just by the coarea formula (see [3]).

The author is indebted to Diogo Oliveira e Silva for suggesting to look at the special exponents under consideration here.

2. Proof of the result

We divide the proof of Theorem 1.3 in two by proving separately two propositions. First we prove

Proposition 2.1. *Let $d \geq 3$. The operator $\mathscr{M}$ satisfies*

$$\|\mathscr{M} f\|_{L^q(\mathbb{S}^{d-1},\mathrm{d}\sigma)} \lesssim_{p,q,d} \|f\|_{L^p(\mathbb{R}^d)} \tag{2.1}$$

for $1 \leq p \leq 4/3$, $p' \geq \frac{d+1}{d-1}q$.

Proof. It suffices to prove the endpoint, that is $p = 4/3$ and $q = 4\frac{d+1}{d-1}$. Let

$$q_d := 4\frac{d+1}{d-1}.$$

By the Stein-Tomas theorem (see e.g. [5]) one has that for the sphere $\mathbb{S}^{d-1}$ immersed in $\mathbb{R}^d$ it holds that for every $f \in L^{4/3}(\mathbb{R}^d)$

$$\|\widehat{f}\|_{L^{q_d}(\mathbb{S}^{d-1},\mathrm{d}\sigma)} \lesssim \|f\|_{L^{4/3}(\mathbb{R}^d)}. \tag{2.2}$$

By duality this is equivalent to the estimate

$$\|\widehat{g\mathrm{d}\sigma}\|_{L^4(\mathbb{R}^d)} \lesssim \|g\|_{L^{q'_d}(\mathbb{S}^{d-1},\mathrm{d}\sigma)}.$$

The numerology here is particularly fortunate since 4 is an even exponent, which allows us to multilinearise and use Plancherel to write

$$\|\widehat{g\mathrm{d}\sigma}\|^2_{L^4(\mathbb{R}^d)} = \|(\widehat{g\mathrm{d}\sigma})^2\|_{L^2(\mathbb{R}^d)} = \|g\mathrm{d}\sigma * g\mathrm{d}\sigma\|_{L^2(\mathbb{R}^d)}$$

(here of course the L^2 norm on the right hand side has to be interpreted as the operator norm of the linear operator given by $h \mapsto \langle g\mathrm{d}\sigma * g\mathrm{d}\sigma, h\rangle$). Thus the Stein-Tomas type estimate (2.2) can be stated equivalently in this case as

$$\|g\mathrm{d}\sigma * g\mathrm{d}\sigma\|_{L^2(\mathbb{R}^d)} \lesssim \|g\|^2_{L^{q'_d}(\mathbb{S}^{d-1},\mathrm{d}\sigma)},$$

which means

$$\Big|\int_{\mathbb{S}^{d-1}}\int_{\mathbb{S}^{d-1}} g(\omega)g(\omega')h(\omega'-\omega)\mathrm{d}\sigma(\omega)\mathrm{d}\sigma(\omega')\Big| \lesssim \|g\|^2_{L^{q'_d}(\mathbb{S}^{d-1},\mathrm{d}\sigma)}\|h\|_{L^2(\mathbb{R}^d)}. \quad (2.3)$$

We linearise the maximal operator $\mathscr{M}$ by defining

$$\mathscr{A}_{\varepsilon(\cdot)}f(\omega) := \int_{\mathbb{R}^d} \widehat{f}(\omega+y)\chi_{\varepsilon(\omega)}(y)\mathrm{d}y,$$

where $\varepsilon(\cdot)$ an arbitrary measurable function that takes positive values. To bound $\mathscr{M}$ it suffices to bound $\mathscr{A}_{\varepsilon(\cdot)}$ in the same range independently of $\varepsilon(\cdot)$. The desired inequality

$$\|\mathscr{A}_{\varepsilon(\cdot)}f\|_{L^{q_d}(\mathbb{S}^{d-1},\mathrm{d}\sigma)} \lesssim \|f\|_{L^{4/3}(\mathbb{R}^d)}$$

is equivalent by duality to the inequality

$$\|\mathscr{A}^*_{\varepsilon(\cdot)}g\|_{L^4(\mathbb{R}^d)} \lesssim \|g\|_{L^{q'_d}(\mathbb{S}^{d-1},\mathrm{d}\sigma)},$$

where $\mathscr{A}^*_{\varepsilon(\cdot)}$ is the formal adjoint of $\mathscr{A}_{\varepsilon(\cdot)}$, which is given by

$$\mathscr{A}^*_{\varepsilon(\cdot)}g(x) := \int_{\mathbb{S}^{d-1}} g(\omega)e^{i\omega\cdot x}\widehat{\chi}(\varepsilon(\omega)x)\mathrm{d}\sigma(\omega).$$

As before, this is equivalent to establishing

$$\|\widehat{\mathscr{A}^*_{\varepsilon(\cdot)}g} * \widehat{\mathscr{A}^*_{\varepsilon(\cdot)}g}\|_{L^2(\mathbb{R}^d)} \lesssim \|g\|^2_{L^{q'_d}(\mathbb{S}^{d-1},\mathrm{d}\sigma)}.$$

First of all, observe that by Fubini's theorem

$$\begin{aligned}\widehat{\mathscr{A}^*_{\varepsilon(\cdot)}g}(\xi) &= \int e^{-ix\cdot\xi}\int_{\mathbb{S}^{d-1}} g(\omega)e^{i\omega\cdot x}\widehat{\chi}(\varepsilon(\omega)x)\mathrm{d}\sigma(\omega)\mathrm{d}x\\ &= \int_{\mathbb{S}^{d-1}} g(\omega)\int e^{-ix\cdot(\xi-\omega)}\widehat{\chi}(\varepsilon(\omega)x)\mathrm{d}x\mathrm{d}\sigma(\omega)\\ &= \int_{\mathbb{S}^{d-1}} g(\omega)\chi_{\varepsilon(\omega)}(\xi-\omega)\mathrm{d}\sigma(\omega)\end{aligned}$$

(with a little abuse of notation). Let then $h \in L^2(\mathbb{R}^d)$, so that by the above observation and multiple applications of Fubini's theorem we have the following

chain of equalities:

$$\begin{aligned}
\langle \widehat{\mathscr{A}^*_{\varepsilon(\cdot)}g} * \widehat{\mathscr{A}^*_{\varepsilon(\cdot)}g}, h\rangle &= \int\int \widehat{\mathscr{A}^*_{\varepsilon(\cdot)}g}(\xi-\eta)\widehat{\mathscr{A}^*_{\varepsilon(\cdot)}g}(\eta)\overline{h(\xi)}\mathrm{d}\eta\mathrm{d}\xi \\
&= \int\int\int_{\mathbb{S}^{d-1}}\int_{\mathbb{S}^{d-1}} g(\omega)g(\omega')\chi_{\varepsilon(\omega)}(\xi-\eta-\omega)\chi_{\varepsilon(\omega')}(\eta-\omega')\mathrm{d}\sigma(\omega)\mathrm{d}\sigma(\omega')\overline{h(\xi)}\mathrm{d}\eta\mathrm{d}\xi \\
&= \int\int_{\mathbb{S}^{d-1}}\int_{\mathbb{S}^{d-1}} g(\omega)g(\omega')\overline{h(\xi)}\Big(\int \chi_{\varepsilon(\omega)}(\xi-\eta-\omega)\chi_{\varepsilon(\omega')}(\eta-\omega')\mathrm{d}\eta\Big)\mathrm{d}\sigma(\omega)\mathrm{d}\sigma(\omega')\mathrm{d}\xi \\
&= \int\int_{\mathbb{S}^{d-1}}\int_{\mathbb{S}^{d-1}} g(\omega)g(\omega')\overline{h(\xi)}(\chi_{\varepsilon(\omega)} * \chi_{\varepsilon(\omega')})(\xi+\omega'-\omega)\mathrm{d}\sigma(\omega)\mathrm{d}\sigma(\omega')\mathrm{d}\xi \\
&= \int_{\mathbb{S}^{d-1}}\int_{\mathbb{S}^{d-1}} g(\omega)g(\omega')\int \tilde{h}(-\xi)(\chi_{\varepsilon(\omega)} * \chi_{\varepsilon(\omega')})(\xi+\omega'-\omega)\mathrm{d}\xi\mathrm{d}\sigma(\omega)\mathrm{d}\sigma(\omega') \\
&= \int_{\mathbb{S}^{d-1}}\int_{\mathbb{S}^{d-1}} g(\omega)g(\omega')(\tilde{h} * \chi_{\varepsilon(\omega)} * \chi_{\varepsilon(\omega')})(\omega'-\omega)\mathrm{d}\sigma(\omega)\mathrm{d}\sigma(\omega'),
\end{aligned}$$

where $\tilde{h}(\xi) = \overline{h(-\xi)}$. But then we have that pointwise

$$|(\tilde{h} * \chi_{\varepsilon(\omega)} * \chi_{\varepsilon(\omega')})(\omega'-\omega)| \lesssim M^2\tilde{h}(\omega'-\omega),$$

with constant depending only on the choice of χ, where M is the Hardy-Littlewood maximal function; therefore by the Stein-Tomas restriction estimate (2.3) we have

$$\begin{aligned}
|\langle \widehat{\mathscr{A}^*_{\varepsilon(\cdot)}g} * \widehat{\mathscr{A}^*_{\varepsilon(\cdot)}g}, h\rangle| &\lesssim \int_{\mathbb{S}^{d-1}}\int_{\mathbb{S}^{d-1}} |g(\omega)||g(\omega')|M^2\tilde{h}(\omega'-\omega)\mathrm{d}\sigma(\omega)\mathrm{d}\sigma(\omega') \\
&\lesssim \|g\|^2_{L^{q'_d}(\mathbb{S}^{d-1},\mathrm{d}\sigma)}\|M^2\tilde{h}\|_{L^2(\mathbb{R}^d)} \lesssim \|g\|^2_{L^{q'_d}(\mathbb{S}^{d-1},\mathrm{d}\sigma)}\|h\|_{L^2(\mathbb{R}^d)},
\end{aligned}$$

which proves the desired estimate for $\mathscr{A}_{\varepsilon(\cdot)}$. □

Remark. It is interesting to notice that the critical endpoint for Fourier restriction to curves in $\mathbb{R}^2$ is $L^{4/3} \to L^{4/3}$, and we know that the corresponding (even restricted) strong type estimate is false by work of Beckner, Carbery, Semmes and Soria [1]. Thus the proof above barely misses to adapt to case $d = 2$.

Finally, we prove the second half of Theorem 1.3, restated below.

Proposition 2.2. *Let $1 \le p \le 8/7$. If $f \in L^p(\mathbb{R}^d)$ then σ-a.e. point of $\mathbb{S}^{d-1}$ is a Lebesgue point for $\widehat{f}$.*

Proof. The proof that follows is taken from [2] and has been included only for the reader's convenience.
Let $\mathcal{R}$ denote the Fourier restriction operator to the sphere $\mathbb{S}^{d-1}$. Let $\mathscr{M}^+$ denote the positive maximal Fourier restriction operator associated to the sphere $\mathbb{S}^{d-1}$, defined as

$$\mathscr{M}^+ f(\omega) := \sup_{\varepsilon>0}\frac{1}{\varepsilon^d}\int_{|y|\le\varepsilon}|\widehat{f}(\omega+y)|\mathrm{d}y.$$

To prove the proposition it suffices to show that

$$\|\mathscr{M}^+ f\|_{L^q(\mathbb{S}^{d-1},\mathrm{d}\sigma)} \lesssim_{p,q} \|f\|_{L^p(\mathbb{R}^d)} \tag{2.4}$$

for $1 \le p \le 8/7$ and $p' \ge q\frac{d+1}{d-1}$. Indeed, assuming this holds, one can define

$$F(\omega) := \limsup_{\varepsilon\to 0}\frac{1}{\varepsilon^d}\int_{|y|\le\varepsilon}|\widehat{f}(\omega+y) - \mathcal{R}f(\omega)|\mathrm{d}y;$$

since $\mathcal{R}\varphi = \widehat{\varphi}|_{\mathbb{S}^{d-1}}$ for any $\varphi \in \mathscr{S}(\mathbb{R}^d)$, we have

$$\begin{aligned}
F(\omega) &\le \limsup_{\varepsilon\to 0}\frac{1}{\varepsilon^d}\int_{|y|\le\varepsilon}|\widehat{f-\varphi}(\omega+y)|\mathrm{d}y + |\mathcal{R}(f-\varphi)(\omega)| \\
&\le \mathscr{M}^+(f-\varphi)(\omega) + |\mathcal{R}(f-\varphi)(\omega)|.
\end{aligned}$$

By the Stein-Tomas estimate and (2.4) it follows then that

$$\|F\|_{L^q(\mathbb{S}^{d-1},\mathrm{d}\sigma)} \lesssim \|f-\varphi\|_{L^p(\mathbb{R}^d)}$$

in the given range, and by taking φ to be an approximant of f in L^p norm we see that $\|F\|_{L^q(\mathbb{S}^{d-1},\mathrm{d}\sigma)} = 0$ or equivalently that $F = 0$ σ-a.e., which proves the proposition. Thus it suffices to prove (2.4), and in particular it suffices to prove it under the assumption that $p' = q\frac{d+1}{d-1}$. This will follow from Proposition 2.1.
Observe that by Hölder's inequality we have

$$\frac{1}{\varepsilon^d}\int_{|y|\leq\varepsilon} |\widehat{f}(\omega+y)|\mathrm{d}y \lesssim \Big(\frac{1}{\varepsilon^d}\int_{|y|\leq\varepsilon} |\widehat{f}(\omega+y)|^2\mathrm{d}y\Big)^{1/2};$$

let then $h := f * \tilde{f}$, so that

$$\widehat{h} = |\widehat{f}|^2,$$

and we have

$$\mathscr{M}^+ f \lesssim (\mathscr{M} h)^{1/2}$$

pointwise. Let s be such that $s \leq 4/3$ and

$$\frac{q}{2}\frac{d+1}{d-1} = s';$$

by Proposition 2.1 we have then

$$\begin{aligned}\|\mathscr{M}^+ f\|_{L^q(\mathbb{S}^{d-1},\mathrm{d}\sigma)} &\lesssim \|\mathscr{M} h\|^{1/2}_{L^{q/2}(\mathbb{S}^{d-1},\mathrm{d}\sigma)} \lesssim \|h\|^{1/2}_{L^s(\mathbb{R}^d)} \\ &\leq \|f\|_{L^p(\mathbb{R}^d)},\end{aligned}$$

where $1 + \frac{1}{s} = \frac{2}{p}$ and the last inequality is an application of Young's inequality. Thus it follows that

$$p' = 2s' = q\frac{d+1}{d-1},$$

as desired. Since $s \leq 4/3$, we see that we can only afford $p \leq 8/7$, and this concludes the proof. □

References

[1] W. Beckner, A. Carbery, S. Semmes, and F. Soria, *A note on restriction of the Fourier transform to spheres*, Bull. London Math. Soc. **21** (1989), no. 4, 394–398. MR 998638

[2] Detlef Müller, Fulvio Ricci, and James Wright, *A maximal restriction theorem and Lebesgue points of functions in $\mathcal{F}(L^p)$*, 2016.

[3] Daniel M. Oberlin, *A uniform Fourier restriction theorem for surfaces in $\mathbb{R}^3$*, Proc. Amer. Math. Soc. **132** (2004), no. 4, 1195–1199. MR 2045437

[4] Per Sjölin, *Fourier multipliers and estimates of the Fourier transform of measures carried by smooth curves in R^2*, Studia Math. **51** (1974), 169–182. MR 0385437

[5] Elias M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series, vol. 43, Princeton University Press, Princeton, NJ, 1993, With the assistance of Timothy S. Murphy, Monographs in Harmonic Analysis, III. MR 1232192

Marco Vitturi, Laboratoire de Mathématiques Jean Leray, 2, rue de la Houssinière BP 92208 44322 Nantes Cedex 3.

E-mail address: marco.vitturi@univ-nantes.fr